



Calculus: Volume of Solids

Definite Integral Notes, Examples, and Formulas related to
Disc/Washer and Shell/Cylinder Methods

Includes Practice Test (with solutions)






Shell (or, Cylinder) Method

Utilizing the Shell (Cylinder) Method

Example: Find the volume of a solid formed by revolving the area bounded by $y = x^2 - 6x + 5$ and $y = 0$ around the x-axis.



Disc Method

$$\text{Volume} = \int_a^b \pi (\text{function})^2 dx$$

$$\text{Volume} = \int_1^5 \pi (x^2 - 6x + 5)^2 dx$$

$$\text{Volume} = \pi \int_1^5 x^4 - 12x^3 + 46x^2 - 60x + 25 dx$$

$$\pi \left(\frac{x^5}{5} - 3x^4 + \frac{46x^3}{3} - 30x^2 + 25x \right) \Big|_1^5$$

$$\boxed{\frac{512}{15} \pi}$$

$$\begin{array}{r} (x^2 - 6x + 5)(x^2 - 6x + 5) \\ x^4 - 6x^3 + 5x^2 \\ -6x^3 + 36x^2 - 30x \\ \hline 5x^2 - 30x + 25 \\ x^4 - 12x^3 + 46x^2 - 60x + 25 \end{array}$$

Now, suppose we want to find the volume of a solid formed from the same area revolved around the y-axis.

If we use the disc method, we need to express the equations in terms of y.



What is the inverse of $y = x^2 - 6x + 5$?

This is difficult to figure out..

However, there is another approach to finding the volume of a solid. the "shell method".

It takes partitions that are parallel to the axis of rotation.

When a partition is revolved around the axis, it creates a cylinder ('shell').



The middle partition revolved around the y-axis forms a cylinder

This lateral area is 24π

(the sum of lateral areas from all the cylinders is the volume of the solid)

Shell (Cylinder) Method

$$\text{Volume} = \int_a^b 2\pi (\text{radius})(\text{height}) dx$$

$$\text{Volume} = 2\pi \int_1^5 (x)(x^2 - 6x + 5) dx$$

$$\text{Volume} = 2\pi \int_1^5 x^3 - 6x^2 + 5x dx$$

$$2\pi \cdot \left(\frac{x^4}{4} - 2x^3 + \frac{5x^2}{2} \right) \Big|_1^5$$

$$2\pi \cdot \left(\frac{624}{4} - 248 + \frac{120}{2} \right) = -64\pi$$

since it's area under the x-axis, the value was negative...

But, area cannot be negative...

$$\boxed{64\pi}$$

Find the volume of a solid formed by revolving the area bounded by

$$y = \sqrt{x} \quad y = 0 \quad \text{and} \quad x = 9$$

around the x-axis

Disc Method

$$\text{Volume} = \int_a^b \pi (\text{function})^2 dx$$

$$\text{Volume} = \int_0^9 \pi (\sqrt{x})^2 dx$$



$$\text{Volume} = \pi \int_0^9 x dx$$

$$\pi \cdot \frac{x^2}{2} \Big|_0^9 = \frac{81}{2} \pi$$

Partitions are perpendicular to the x-axis

Each partition is a disc!

(the radius of each disc is the function)



Volume of Solids: Shell vs. Disc Method

Lateral Area of cylinder:

$$LA = 2\pi(\text{radius})(\text{height})$$

Shell Method

$$\text{Volume} = \int_a^b 2\pi (\text{radius})(\text{height}) dy$$

$$y = \sqrt{x} \longrightarrow x = y^2$$

$$\text{Volume} = \int_0^3 2\pi (y)(y^2) dy$$


$$\text{Volume} = 2\pi \int_0^3 (y^3) dy$$

$$2\pi \cdot \frac{y^4}{4} \Big|_0^3 = \frac{81}{2} \pi$$


Partitions are parallel to the x-axis

Each partition is a cylinder!

(the height of each cylinder is the function, and the radius of each cylinder is y)



front view of partition



side view of partition
(hollow cylinder)



Each cylinder is a 'shell'.

When all the different sized shells are added together, they form the solid.

(Depending on the function, orientation, and/or rotation, one method may be easier than the other....)

Example: What is the volume of the solid from the region between the x-axis and $y = -x^2 + 4x - 3$ revolved around the y-axis?

Method 1: Apply the "cylinder method" (or "shell method")

Note: each partition is a cylinder with

radius: x

height: $-x^2 + 4x - 3$

formula for surface area of cylinder:

$$SA = 2\pi(\text{radius})(\text{height})$$


We'll construct a definite integral that represents cylinder partitions from $x = 1$ to 3

$$\int_1^3 2\pi x (-x^2 + 4x - 3) dx$$

(radius) (height)

$$2\pi \int_1^3 -x^3 + 4x^2 - 3x dx \Rightarrow 2\pi \left[-\frac{x^4}{4} + \frac{4x^3}{3} - \frac{3x^2}{2} \right]_1^3 =$$

$$2\pi \left(-\frac{81}{4} + 36 - \frac{27}{2} + \frac{1}{4} - \frac{4}{3} + \frac{3}{2} \right) = \boxed{\frac{16}{3}\pi}$$



Method 2: Using "Disk Method" (and horizontal partitions)

Since we're using horizontal partitions, we need to solve for x

$$y = -x^2 + 4x - 3 \quad \text{solve for } x:$$

$$y + 3 = -x^2 + 4x$$

$$y + 3 = -1(x^2 - 4x)$$

(complete the square)

$$-4 + y + 3 = -1(x^2 - 4x + 4)$$

$$y - 1 = -1(x - 2)^2$$

$$1 - y = (x - 2)^2$$

$$\pm \sqrt{1 - y} = (x - 2)$$

$$x = 2 \pm \sqrt{1 - y}$$

right half


left half

$$x = 2 + \sqrt{1 - y} \quad x = 2 - \sqrt{1 - y}$$

Note: each partition is a "washer"

Outer disk: radius extends from y-axis to right half of curve

Inner disk (that is cut out): radius extends from y-axis to left half of curve



We'll construct definite integrals that represent the horizontal disk partitions extending from $y = 0$ to $y = 1$

$$\int_0^1 \pi (2 + \sqrt{1 - y})^2 dy - \int_0^1 \pi (2 - \sqrt{1 - y})^2 dy$$

(radius of 'outer' disk) (radius of 'inner' disk that is being hollowed out)


$$\pi \int_0^1 4 + 4\sqrt{1 - y} + (1 - y) dy - \pi \int_0^1 4 - 4\sqrt{1 - y} + (1 - y) dy$$

$$\pi \int_0^1 5 + 4\sqrt{1 - y} - y dy - \pi \int_0^1 5 - 4\sqrt{1 - y} - y dy$$

$$\frac{43}{6}\pi - \frac{11}{6}\pi = \boxed{\frac{16}{3}\pi}$$

$$\pi \int_0^1 8\sqrt{1 - y} dy$$

$$8\pi \left[\frac{-(1 - y)^{3/2}}{3/2} \right]_0^1 = \boxed{\frac{16}{3}\pi}$$



The Math Guy enjoys his new profession...



Practice Test →

Volume of Solids Quiz: Disc and Shell Methods

1) Find the volume of a solid with area bounded by $y = x^2$ and $y = 4x - x^2$, and

a) revolved around the y-axis

b) revolved around $x = 4$

2) Find the volume of a solid with area bounded by

$$y = x^3 \quad y = 0 \quad \text{and} \quad x = 2$$

and revolved around:

a) x-axis

b) y-axis

c) the line $x = 4$

d) the line $y = 8$

- 3) The region R is bounded by $y = \sqrt{x}$
 $x = 0$
 $y = 2$


Find the volume of the region R rotated around $x = 4$,


- a) using the disk method
- b) using the shell method

- 4) Find the volume of the solid generated by revolving the ellipse $9x^2 + 16y^2 = 144$

- a) around its major axis
- b) around its minor axis

Use disk and shell methods...





Happy Days

*Despite Richie's help, Fonzie dropped out of Calculus.
(... although he did have some success with velocity and acceleration!)*

Solutions-→

1) Find the volume of a solid with area bounded by $y = x^2$ and $y = 4x - x^2$, and

a) revolved around the y-axis

b) revolved around $x = 4$

Since it is difficult to put the second equation in terms of y , we'll utilize the 'shell method' (and use parallel partitions)

$$\text{a) Volume} = \int_a^b 2\pi(\text{radius})(\text{height}) \, dx$$

$$\text{Volume} = \int_0^2 2\pi(x)((4x - x^2) - x^2) \, dx$$

$$\begin{aligned} \text{Volume} &= 2\pi \int_0^2 (4x^2 - 2x^3) \, dx \\ &= 2\pi \left(\frac{4x^3}{3} - \frac{x^4}{2} \right) \Big|_0^2 = 2\pi \left(\frac{32}{3} - 8 \right) = \frac{16\pi}{3} \end{aligned}$$

b) the 'radius': $(4 - x)$


the 'height': $(4x - x^2) - x^2$

$$\text{Volume} = 2\pi \int_0^2 (4-x)(4x - 2x^2) \, dx$$

radius & height
of each shell/cylinder


$$\begin{aligned} \text{Volume} &= 2\pi \int_0^2 (16x - 8x^2 - 4x^2 + 2x^3) \, dx \\ &= 2\pi \left(8x^2 - 4x^3 + \frac{x^4}{2} \right) \Big|_0^2 \end{aligned}$$

$$2\pi (32 + 32 + 8 - 0 + 0 - 0) = 16\pi$$



Observe: Each cylinder (shell) has radius x and height $(4x - x^2) - x^2$

and, $2\pi(\text{radius})(\text{height})$ is the surface area of each cylinder!



2) Find the volume of a solid with area bounded by

$$y = x^3 \quad y = 0 \quad \text{and} \quad x = 2$$

and revolved around:


- a) x-axis
- b) y-axis
- c) the line $x = 4$
- d) the line $y = 8$

a) Utilizing the disc method:
x-axis

$$\text{Volume} = \int_a^b \pi (\text{radius})^2 dx$$

$$\text{Volume} = \int_0^2 \pi (x^3)^2 dx$$

$$\pi \left. \frac{x^7}{7} \right|_0^2 = \boxed{\frac{128\pi}{7}}$$



b) Utilizing the shell method:
y-axis

$$\text{Volume} = \int_a^b 2\pi (\text{radius})(\text{height}) dx$$

$$\text{Volume} = 2\pi \int_0^2 (x)(x^3) dx$$

$$2\pi \left. \frac{x^5}{5} \right|_0^2 = \boxed{\frac{64\pi}{5}}$$


c) Using the shell (cylinder) method:
the line $x = 4$

$$\text{Volume} = \int_0^2 2\pi (4-x)(x^3) dx$$

radius
height

$$\text{Volume} = 2\pi \int_0^2 (4x^3 - x^4) dx$$

$$2\pi \left(x^4 - \frac{x^5}{5} \right) \Big|_0^2 = 32\pi - \frac{64\pi}{5} = \boxed{\frac{96\pi}{5}}$$



d) Using the shell method:
the line $y = 8$

$$\text{Volume} = 2\pi \int_0^8 (8-y)(2 - \sqrt[3]{y}) dy$$

$$\text{Volume} = 2\pi \int_0^8 (16 - 8y^{1/3} - 2y + y^{4/3}) dy$$

$$2\pi \left(16y - \frac{4}{3}y^{4/3} - y^2 + \frac{3y^{7/3}}{7} \right) \Big|_0^8$$

$$2\pi (128 - 96 - 64 + \frac{384}{7}) = \boxed{\frac{320\pi}{7}}$$

d) Using the disk / washer method:

$$\text{Volume} = \int_0^2 \pi (8)^2 - \pi (8 - x^3)^2 dx$$

$$\text{Volume} = \pi \int_0^2 (64 - 64 - 16x^3 + x^6) dx$$

$$\pi \left(-4x^4 + \frac{x^7}{7} \right) \Big|_0^2 = -64 + \frac{128}{7}$$

It's negative because the region is below $y = 8$. Of course, area must be positive...

$$\boxed{-\frac{320}{7}\pi}$$

3) The region R is bounded by $y = \sqrt{x}$
 $x = 0$
 $y = 2$

SOLUTIONS

Find the volume of the region R rotated around $x = 4$,

- a) using the disk method
- b) using the shell method



DISK $\pi(\text{radius})^2$ (area of a disc/circle)

$$\pi \int_0^2 (4)^2 dy - \pi \int_0^2 (4 - y^2)^2 dy$$

radius of outside radius of inside

$$\pi \left[16y \right]_0^2 - \pi \left(16y - \frac{8y^3}{3} + \frac{y^5}{5} \right) \Big|_0^2$$

$$32\pi - \left(32 - \frac{64}{3} + \frac{32}{5} \right) \pi = \frac{224\pi}{15}$$



each partition is a flat disc with the center hallowed out...

SHELL

$2\pi(\text{radius})(\text{height})$ (lateral area of a cylinder)


$$\int_0^4 2\pi(4-x)(2-\sqrt{x}) dx$$

radius height

$$2\pi \int_0^4 8 - 2x - 4\sqrt{x} + \frac{3}{2}x^2 dx$$

$$2\pi \left(8x - x^2 - \frac{8x^{3/2}}{3} + \frac{1}{2}x^2 \right) \Big|_0^4$$

$$2\pi \left(32 - 16 - \frac{64}{3} + \frac{64}{5} \right) = 2\pi \left(16 - \frac{128}{15} \right) = \frac{224\pi}{15}$$



each partition is a cylinder shell

4) Find the volume of the solid generated by revolving the ellipse $9x^2 + 16y^2 = 144$

SOLUTIONS


a) around its major axis

Use disk and shell methods...

b) around its minor axis

In standard form:

$$\frac{x^2}{16} + \frac{y^2}{9} = 1$$



a) major axis is the x-axis

Since the ellipse (ellipsoid) is symmetric, we'll focus on the top 1/2 of the figure...

Then, using the disk method...

$$\frac{y^2}{9} = 1 - \frac{x^2}{16}$$

$$y^2 = 9 - \frac{9x^2}{16}$$

each radius is

$$y = \pm \sqrt{9 - \frac{9x^2}{16}}$$

The boundaries are -4 to 4

$$\int_{-4}^4 \pi (\text{radius})^2 dx$$

$$\int_{-4}^4 \pi \left(9 - \frac{9x^2}{16} \right) dx = \pi \left(9x - \frac{3x^3}{16} \right) \Big|_{-4}^4 = \pi \left((36 - 12) - (-36 + 12) \right) = 48\pi$$

b) minor axis is the y-axis

Again, the ellipse (and ellipsoid) is symmetric over the axis, so we can work on half.

Then, using the shell method...

$$\frac{y^2}{9} = 1 - \frac{x^2}{16}$$

$$y^2 = 9 - \frac{9x^2}{16}$$


$$y = \pm \sqrt{9 - \frac{9x^2}{16}}$$

The cylinders/shells have vertical heights, and we'll focus on quadrant I... So the boundaries will be 0 to 4.

$$\int_0^4 2\pi (\text{radius})(\text{height})$$

$$\int_0^4 2\pi x \sqrt{9 - \frac{9x^2}{16}} dx$$

Radii Height



Using a calculator ----> the integral is 32π


Finally, we double the answer to account for the bottom half of the ellipsoid!

$$64\pi$$

Thanks for visiting. (Hope it helped!)

If you have questions, suggestions, or requests, let us know.

Cheers



Also, mathplane *express* for mobile and tablets at mathplane.org

And, facebook, TeachersPayTeachers, TES and Pinterest